

A CLASS OF COUNTER-EXAMPLES TO THE HYPERSECTION PROBLEM BASED ON FORCING EQUATIONS

HOLGER BRENNER

ABSTRACT. We give a class of three-dimensional Stein spaces W together with a hypersurface H , such that the complement $W - H$ is not Stein, but such that for every analytic surface $S \subset W$ the complement $S - S \cap H$ is Stein. This class is constructed using forcing equations and gives new counter-examples to the hypersection problem.

Mathematical Subject Classification (1991): 14J26, 32C25, 32L05, 32E10

0. Introduction. Let W be a complex Stein space of dimension ≥ 3 and let $H \subset W$ be an analytic hypersurface, $U = W - H$. Suppose that for every analytic hypersurface $S \subset W$ the intersection $U \cap S$ is Stein, is then U itself Stein? This question is called the hypersection problem, see [2] for a general treatment and related problems. The first counter-example to this question was given by Coltoiu and Diederich in [1], using the affine cone over the complement of two sections on some ruled surface over an elliptic curve. In this way they get a normal three-dimensional isolated singularity.

In this paper we present another class of three-dimensional Stein spaces W together with a hypersurface H fulfilling the assumptions in the hypersection problem, but not its conclusion. The class is constructed in the following way: we start with a two-dimensional normal affine cone X over a smooth projective curve and the vertex point $P \in X$. Suppose that we have three homogeneous functions f_1, f_2 and f_0 on X . Then, under suitable conditions, $W = V(f_1 t_1 + f_2 t_2 + f_0) \subset X \times \mathbb{C}^2$ and the hypersurface $H = p^{-1}(P)$ have the desired properties, see Theorem 7. These conditions reduce to numerical conditions, which are easily to verify, see Corollary 8 and Example 9.

1. Forcing equations. Let X be an irreducible normal complex Stein space of dimension d together with a point $P \in X$. Let $f_1, \dots, f_n$ be holomorphic functions on X such that the common zero set of these functions is exactly P . Let f_0 be another holomorphic function vanishing at P . Then we consider the complex space $W \subset X \times \mathbb{C}^n$ defined by the equation $f_1 t_1 + \dots + f_n t_n + f_0 = 0$,

$$W = \{(x, t_1, \dots, t_n) \in X \times \mathbb{C}^n : f_1(x)t_1 + \dots + f_n(x)t_n + f_0(x) = 0\}.$$

The equation $f_1 t_1 + \dots + f_n t_n + f_0 = 0$ is called a forcing equation, since it forces f_0 to lie in the ideal generated by $f_1, \dots, f_n$. Forcing equations and the algebras defined by them play an important role in the theory of closure operations for ideals, e.g. tight closure and solid closure, see [5]. Let $p : W \rightarrow X$ be the projection. If $d \geq 2$, then $n \geq 2$ and W is an irreducible Stein space of dimension $d + n - 1$. Let $U_i = \{x \in X : f_i(x) \neq 0\}$ and $U = \bigcup_{i=1}^n U_i = X - P$. Resolving t_i shows that $p^{-1}(U_i) \cong U_i \times \mathbb{C}^{n-1}$. The transition mappings however are only affine-linear, not

linear, hence $W|_U$ is not a vector bundle. $H := p^{-1}(P) \cong \mathbb{C}^n$ is a closed subset in W , which is a hypersurface in case $d = 2$. The existence of a section $X \rightarrow W$ is equivalent with $f_0 \in (f_1, \dots, f_n)$ over X .

Lemma 1. *Let X be a normal irreducible Stein space together with a point P and let $f_1, \dots, f_n, f_0 \in \Gamma(X, \mathcal{O}_X)$ be holomorphic functions on X . Let X' be another irreducible Stein space of the same dimension and let $\psi : X' \rightarrow X$ be a holomorphic mapping such that $\psi^{-1}(P)$ contains isolated points. Suppose that $f_0 \circ \psi \in (f_1 \circ \psi, \dots, f_n \circ \psi)$ in $\Gamma(X', \mathcal{O}_{X'})$. Then $f_0 \in (f_1, \dots, f_n)$ in $\mathcal{O}_{X,P}$.*

Proof. Let Q be an isolated point over P . Then there exist open neighborhoods $Q \in U$ and $P \in V$ such that $\psi : U \rightarrow V$ is finite, see [4], Ch. 3.2. Due to the finite mapping theorem, $\psi_*(\mathcal{O}_{X'})$ is a coherent analytic algebra on $P \in V \subseteq X$. Furthermore it is torsionfree due to the assumptions on the dimension. Since X is normal, we have the trace map $tr : \psi_*(\mathcal{O}_U) \rightarrow \mathcal{O}_V$, which gives the result. $\square$

Corollary 2. *Let X be a normal irreducible Stein space of dimension d together with a point P and let $f_1, \dots, f_n, f_0 \in \Gamma(X, \mathcal{O}_X)$ be holomorphic functions such that $f_0 \notin (f_1, \dots, f_n)\mathcal{O}_P$. Let*

$$W = V(f_1 t_1 + \dots + f_n t_n + f_0) \subset X \times \mathbb{C}^n \xrightarrow{p} X.$$

and $H = p^{-1}(P)$. Let T be an irreducible complex space of dimension d and let $\varphi : T \rightarrow W$ be a holomorphic map. Then $\varphi^{-1}(H) \subset T$ contains no isolated points and the codimension of $\varphi^{-1}(H)$ is $\leq d - 1$.

Proof. We look at the composed mapping $\psi = p \circ \varphi : T \xrightarrow{\varphi} W \xrightarrow{p} X$. Since it factors through W it follows that $f_0 \circ \psi \in (f_1 \circ \psi, \dots, f_n \circ \psi)$ in $\Gamma(T, \mathcal{O}_T)$. Due to the Lemma $\psi^{-1}(P) = \varphi^{-1}(H)$ cannot contain isolated points. $\square$

With this result we can establish the hypothesis of the hypersection problem in a broad class of example where the base space X is two-dimensional, $n = 2$ and $H = p^{-1}(P) \cong \mathbb{C}^2$ is a hypersurface in three-dimensional W .

Proposition 3. *Let X be a normal irreducible two-dimensional Stein space together with a point P . Let $f_1, f_2, f_0 \in \Gamma(X, \mathcal{O}_X)$ be holomorphic functions such that $P = \{f_1 = f_2 = 0\}$ and suppose that $f_0 \notin (f_1, f_2)\mathcal{O}_P$. Let*

$$W = Z(f_1 t_1 + f_2 t_2 + f_0) \subset X \times \mathbb{C}^2 \xrightarrow{p} X.$$

and let $H = p^{-1}(P) \subset W$. Then for every analytic surface $S \subset W$ the complement of $S \cap H \subset S$ is Stein.

Proof. We may assume that S is irreducible, let $\tilde{S}$ be its normalization and let $\varphi : \tilde{S} \rightarrow W$ be the corresponding mapping. Due to Cor. 2 we know that $\varphi^{-1}(H) \subset \tilde{S}$ contains no isolated points. Hence $\varphi^{-1}(H)$ is a pure curve on a normal Stein surface and due to [7] its complement is Stein. But then also $S - S \cap H$ itself is Stein. $\square$

2. The graded situation. We have to look for examples of the type described in Proposition 3 where $W - H$ is not Stein. To this end we look at the graded situation. Let $Y \subseteq \mathbb{P}^N$ be a smooth projective variety with the very ample line bundle $H_Y \rightarrow Y$ (which is the restriction of $\mathcal{O}_{\mathbb{P}^N}(1)$ to Y) and let $X \subseteq \mathbb{C}^{N+1}$ be the corresponding affine cone. Let P be the vertex of the cone and assume that X is normal. Recall that we have an action of $\mathbb{C}^*$ on X , which is free on $U = X - P$.

Y is the quotient of this action and $U \rightarrow Y$ is a $\mathbb{C}^*$ -principal bundle. A number $e \in \mathbb{Z}$ defines the action on $X \times \mathbb{C}$ by $\lambda(x, t) := (\lambda x, \lambda^e t)$, this action is free over U and the quotient is the line bundle $H_Y^e \rightarrow Y$.

Suppose that the holomorphic functions f_i are homogeneous of degree d_i , i.e. $f_i(\lambda x) = \lambda^{d_i} f_i(x)$, $x \in X$, $\lambda \in \mathbb{C}^*$. We may consider a homogeneous holomorphic function f of degree d as a section $Y \rightarrow H_Y^d$ and as a mapping of line bundles $H_Y^{-d} \rightarrow Y \times \mathbb{C}$ or $H_Y^e \rightarrow H_Y^{e+d}$.

Proposition 4. *Let X be a normal affine cone over a smooth projective variety Y and let P be the vertex point. Let $f_1, \dots, f_n$ be homogeneous functions such that $P = \{f_1 = \dots = f_n = 0\}$, $U = X - P$. Let d_i be the degrees of f_i and let e_i numbers such that $m = d_i + e_i$ is constant. Then the following hold.*

(i) *There is an exact sequence of vector bundles*

$$0 \longrightarrow V_m \longrightarrow H_Y^{e_1} \times_Y \dots \times_Y H_Y^{e_n} \xrightarrow{\sum f_i} H_Y^m \longrightarrow 0.$$

- (ii) V_m is the quotient of $V(f_1 t_1 + \dots + f_n t_n)|_U$ by the action of $\mathbb{C}^*$ given by $\lambda(x, t_1, \dots, t_n) = (\lambda x, \lambda^{e_1} t_1, \dots, \lambda^{e_n} t_n)$.
- (iii) We have $\text{Det } V_m \cong H_Y^k$, where $k = \sum_{i=1}^n e_i - m = -\sum_{i=1}^n d_i + (n-1)m$.
- (iv) $V_{m'} = V_m \otimes H_Y^{(m'-m)}$. The projective bundle $\mathbb{P}(V_m)$ is independent of the choosen degree m .

Proof. (i). We consider the f_i as morphisms $H_Y^{e_i} \rightarrow H_Y^m$. The morphism of vector bundles $H_Y^{e_1} \times_Y \dots \times_Y H_Y^{e_n} \xrightarrow{\sum f_i} H_Y^m$ over Y is surjective, because the f_i do not have a common zero on Y . Hence the kernel is a vector bundle V_m on Y of rank $n-1$.

(ii). The pull back under $q : U \rightarrow Y$ of the exact sequence in (i) gives

$$0 \longrightarrow q^* V_m \longrightarrow U \times \mathbb{C}^n \xrightarrow{\sum f_i} U \times \mathbb{C} \longrightarrow 0$$

together with the described action, and $q^* V_m = V(f_1 t_1 + \dots + f_n t_n)|_U$. (iii) and (iv) follow. $\square$

Proposition 5. *Let X be a normal affine cone over a smooth projective variety Y and let P be the vertex point, $U = X - P$. Let $f_1, \dots, f_n$ be homogeneous functions such that $P = \{f_1 = \dots = f_n = 0\}$. Let f_0 be another homogeneous function, $d_i = \deg(f_i)$, and let e_i be numbers such that $m = d_i + e_i$ is constant for $i = 0, \dots, n$. Let V_m (resp. V'_m) be the vector bundle on Y defined in Proposition 4 with respect to $f_1, \dots, f_n$ (resp. $f_0, \dots, f_n$). Then the following hold.*

(i) *There is an exact sequence of vector bundles on Y :*

$$0 \longrightarrow V_m \longrightarrow V'_m \xrightarrow{t_0} H_Y^{e_0} \longrightarrow 0.$$

- (ii) *The corresponding embedding $\mathbb{P}(V_m) \hookrightarrow \mathbb{P}(V'_m)$ is independent of m , $\mathbb{P}(V_m)$ is a divisor on $\mathbb{P}(V'_m)$.*
- (iii) *Let $e_0 = 0$. The normal bundle for $\mathbb{P}(V) \hookrightarrow \mathbb{P}(V')$ on $\mathbb{P}(V)$ is $H_{\mathbb{P}(V)}$, where $H_{\mathbb{P}(V)}$ denotes the relative very ample line bundle on $\mathbb{P}(V)$.*
- (iv) *Let $e_0 = 0$. $W|_U \rightarrow \mathbb{P}(V') - \mathbb{P}(V)$ is a quotient of the action on $W = V(f_1 t_1 + \dots + f_n t_n + f_0)$ given by*

$$\lambda(x, t_1, \dots, t_n) \mapsto (\lambda x, \lambda^{e_1} t_1, \dots, \lambda^{e_n} t_n).$$

Proof. (i). The mappings in the sequence follow from the defining sequences for V_m and V'_m . The exactness of the sequence follows from diagram chasing. (ii) is clear.

(iii). Since we assume $e_0 = 0$, t_0 is a global function on V' and it is a global section in the relative very ample line bundle $H_{\mathbb{P}(V')}$ on $\mathbb{P}(V')$, and $\mathbb{P}(V)$ is the corresponding divisor. Therefore the normal bundle of this embedding is $i^*H_{\mathbb{P}(V')} \cong H_{\mathbb{P}(V)}$.

(iv). First we may identify the closed subset $\{Q \in V' : t_0(Q) = 1\}$ with $\mathbb{P}(V') - \mathbb{P}(V)$. The described action on W respects the forcing equation, for $f_1(\lambda x)\lambda^{e_1}t_1 + \dots + f_n(\lambda x)\lambda^{e_n}t_n + f_0(\lambda x) = \lambda^{d_1}\lambda^{e_1}f_1(x)t_1 + \dots + \lambda^{d_n}\lambda^{e_n}f_n(x)t_n + \lambda^{d_0}f_0(x) = \lambda^m(f_1(x)t_1 + \dots + f_n(x)t_n + f_0(x)) = 0$. This action on $W|_U = W - p^{-1}(P)$ is the same action as the action on the vector bundle $V(f_1t_1 + \dots + f_0t_0)|_U$ restricted to $t_0 = 1$ described in Proposition 4(ii). Its quotient is $\{Q \in V' : t_0(Q) = 1\}$. $\square$

Now we specialize to the two-dimensional situation.

Corollary 6. *Let X be a normal affine two-dimensional cone over a smooth projective curve Y and let P be the vertex point. Let f_1, f_2 be homogeneous functions such that $P = \{f_1 = f_2 = 0\}$. Let f_0 be another homogeneous function, $d_i = \deg(f_i)$, and let e_i numbers such that $m = d_i + e_i$ is constant for $i = 0, 1, 2$. Let V_m (V'_m) be the corresponding vector bundles on Y . Then the following hold.*

- (i) $\mathbb{P}(V')$ is a ruled surface and $\mathbb{P}(V) \subset \mathbb{P}(V')$ is a section (independent of m).
- (ii) We have $V_m \cong H_Y^{e_0+d_0-d_1-d_2}$ and the exact sequence

$$0 \longrightarrow H_Y^{e_0+d_0-d_1-d_2} \longrightarrow V' \longrightarrow H_Y^{e_0} \longrightarrow 0.$$

- (iii) Let $e_0 = 0$. The normal bundle of the embedding $Y \cong \mathbb{P}(V) \subset \mathbb{P}(V')$ is $H_Y^{d_1+d_2-d_0}$.
- (iv) The self intersection number of $Y \cong \mathbb{P}(V) \hookrightarrow \mathbb{P}(V')$ is $(d_1 + d_2 - d_0) \deg H_Y$.

Proof. (i) is clear due to Proposition 5.

(ii) From the defining sequence in Proposition 4 it follows that we have $V_m \cong H_Y^{e_1} \otimes H_Y^{e_2} \otimes H_Y^{-m} = H_Y^{e_1+e_2-m} = H_Y^{e_0+d_0-d_1-d_2}$.

(iii). The normal bundle on $\mathbb{P}(V)$ is $H_{\mathbb{P}(V)}$ due to Proposition 5. But for a line bundle this is just the negative tautological bundle $-V$, therefore $N = -V = H^{d_1+d_2-d_0}$.

(iv). The self intersection number is $\mathbb{P}(V)^2 = \deg_Y N = \deg_Y H^{d_1+d_2-d_0} = (d_1 + d_2 - d_0) \deg H_Y$. $\square$

3. A class of examples.

Theorem 7. *Let X be a normal affine two-dimensional cone with vertex point P over a smooth projective curve Y , let f_1, f_2 and f_0 be homogeneous holomorphic functions on X with degrees d_1, d_2, d_0 such that*

$$(1) V(f_1, f_2) = P, \quad (2) f_0 \notin (f_1, f_2)\mathcal{O}_{X,P} \quad \text{and} \quad (3) d_1 + d_2 - d_0 < 0.$$

Let $W = V(f_1t_1 + f_2t_2 + f_0) \subset X \times \mathbb{C}^2$ and $H = p^{-1}(P) \subset W$. Then $W - H \subset W$ is not Stein, but for every analytic surface $S \subset W$ the intersection $(W - H) \cap S$ is Stein.

Proof. We have to show that $W - H$ is not Stein. Since $W - H = W|_U$, $U = X - P$, the quotient of this open subset under the action of $\mathbb{C}^*$ is $\mathbb{P}(V') - \mathbb{P}(V)$. Due to [6] it is enough to show that this complement of the section in the ruled surface $\mathbb{P}(V')$

is not Stein. The self intersection of $\mathbb{P}(V)$ is $(d_0 + d_1 - d_0) \deg H_Y < 0$, hence $\mathbb{P}(V)$ is due to [3] contractible and the complement cannot be Stein. $\square$

Corollary 8. *Let h be an irreducible homogeneous polynomial of degree r in the three variables x, y, z and suppose that x, y are homogeneous parameters (i. e. that $h = cz^r + \text{other terms}$, $c \neq 0$) and let $X = V(h) \subset \mathbb{C}^3$. Suppose that X is normal (hence Y is smooth). Let $d_1, d_2 \geq 1$ and d_0 be degrees such that $d_1 + d_2 < d_0 < r$. Then $f_1 = x^{d_1}$, $f_2 = y^{d_2}$ and $f_0 = z^{d_0}$ fulfill the conditions of the theorem.*

Proof. We have to show that $z^{d_0} \notin (x^{d_1}, y^{d_2})\mathcal{O}_P$. For this we look at the completion of the local ring, $\hat{\mathcal{O}}_P/(x^{d_1}, y^{d_2}) = \mathbb{C}[[x, y, z]]/(h, x^{d_1}, y^{d_2})$. Since $d_0 < r = \deg(h)$ we see that $z^{d_0} \neq 0$ in this residue class ring. On the other hand, the self intersection number is $(d_1 + d_2 - d_0) \deg H_Y < 0$. $\square$

Example 9. Let $X = V(x^r + y^r + z^r) \subset \mathbb{C}^3$, $r \geq 4$ be a Fermat type hypersurface, let

$$W = V(x^r + y^r + z^r, xt_1 + yt_2 + z^s) \subset \mathbb{C}^5, \quad 3 \leq s < r \quad \text{and} \quad H = V(x, y).$$

Then the conditions in the corollary are fulfilled.

The easiest example of this type is the Fermat quartic $x^4 + y^4 + z^4$ together with $f_1 = x$, $f_2 = y$ and $f_0 = z^3$. Therefore

$$W = V(x^4 + y^4 + z^4, xt_1 + yt_2 + z^3) \quad \text{and} \quad H = V(x, y)$$

gives an counter-example to the hypersection problem.

Remarks 10. The hypersurface H in our example is the singular locus of W . Since the normalization does not change the complement of H and since its preimage is still a hypersurface due to Corollary 2, we also may get normal examples.

The condition $f_0 \notin (f_1, f_2)\mathcal{O}_P$ in Corollary 8 ensures that the divisor $\mathbb{P}(V) \subset \mathbb{P}(V')$ intersects every curve $C \subset \mathbb{P}(V')$ positively. For a disjoint curve would yield a closed punctured surface (its cone) inside $W - H$. If additionally $d_1 + d_2 - d_0 > 0$, then $\mathbb{P}(V)$ is an ample divisor and its complement is affine, hence Stein. What happens if $d_1 + d_2 - d_0 = 0$? Then the complement is not affine, but it may be Stein. For $h = x^3 + y^3 + z^3 = 0$, $f_1 = x$, $f_2 = y$ and $f_0 = z^2$ we get an instance of the classical construction of Serre of a Stein, but non-affine variety, see [8].

REFERENCES

- [1] Coltoiu, M., Diederich, K.: Open sets with Stein hypersurface sections in Stein spaces. *Ann. Math.* **145** (1997), 175-182.
- [2] Diederich, K.: Some Aspects of the Levi Problem: Recent Developments. In *Geometric Complex Analysis* (edited by Junjirō Noguchi et. al.) (1996), 163-181.
- [3] Grauert, H.: Über Modifikationen und exzeptionelle analytische Mengen. *Math. Ann.* **146** (1962), 331-368.
- [4] Grauert, H., Remmert, R.: *Coherent Analytic Sheaves*. Springer 1984.
- [5] Hochster, M.: Solid closure. *Contemp. Math.* **159** (1994), 103-172.
- [6] Matsushima, Y., Morimoto, A.: Sur certain espaces fibrés holomorphes sur une variété de Stein. *Bull. Soc. Math. France* **88** (1960), 127-155.
- [7] Simha, R.R.: On the complement of a curve on a Stein space of dimension two. *Math. Z.* **82** (1963), 63-66.
- [8] Umemura, H.: La dimension cohomologique des surfaces algébriques. *Nagoya Math. J.* **47** (1972), 155-160.

MATHEMATISCHE FAKULTÄT, RUHR-UNIVERSITÄT, UNIVERSITÄTSSTR. 150, 44780 BOCHUM,
GERMANY

E-mail address: `brenner@cobra.ruhr-uni-bochum.de`